

Question solution 2066(2nd batch)

Group A(2X10=20)

Q1)Find the length of curve $y = x^{3/2}$ from $x=0$ to $x=4$.

Solution:

$$y = x^{3/2}$$

$$-----0 \leq x \leq 4$$

$$\frac{dy}{dx} = \frac{3}{2}x^{1/2}$$

$$\frac{dy}{dx} = \frac{9}{4}x$$

$$L = \int_0^4 \sqrt{1 + \frac{9}{4}x} dx$$

$$= \int_4^0 \left(1 + \frac{9}{4}x\right)^{1/2} dx$$

$$= \left[\frac{\left(1 + \frac{9}{4}x\right)^{3/2}}{\frac{3}{2} \times 9/4} \right]$$

$$= \frac{8}{27} \left[\left(1 + 9\right)^{3/2} - 1 \right]$$

$$= \frac{8}{27} \left[(10)^{3/2} - 1 \right] \text{Ans}$$

Q.2) Find critical points of the function $f(x) = x^{3/2}(x-4)$

Solution:

$$F(x) = x^{3/2}(x-4)$$

$$=x^{3/2+1}-4x^{3/2}$$

$$=x^{5/2}-4x^{3/2}$$

$$f'(x)=\frac{5}{2}x^{\frac{3}{2}}-4*\frac{3}{2}x^{\frac{1}{2}}$$

$$=\frac{5}{2}x^{\frac{3}{2}}-\frac{12}{2}x^{\frac{1}{2}}$$

$$=\frac{5}{2}x^{\frac{3}{2}}-6x^{\frac{1}{2}}$$

$$f'(x)=0$$

$$\text{or; } \frac{5}{2}x^{\frac{3}{2}}-6x^{\frac{1}{2}}=0$$

$$\text{or; } \frac{5}{2}x^{\frac{1}{2}}-x^1-6x^{\frac{1}{2}}=0$$

$$\text{or; } x^{\frac{1}{2}}\left[\frac{5}{2}x-6\right]=0$$

$$\text{or; } \frac{5}{2}x-6=0$$

$$\text{or; } x=\frac{12}{5}$$

therefore critical point=(12,0)

Q.N.3) Does the following series Converge?

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

Solution: we have the given series

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\text{Let } f(n)=1/n^2$$

$$\begin{aligned}
 \sum_{n=1}^{\infty} \frac{1}{n^2} &= f(n) = f(0) + f(1) + f(2) + f(3) + \dots f(n) \\
 &= f(1) + \int_1^{\infty} f(n) \\
 &= f(1) + \lim_{b \rightarrow \infty} \int_1^b f(n) \\
 &= f(1) + \lim_{b \rightarrow \infty} \int_1^b \frac{1}{n^2} dx \\
 &= f(1) + \lim_{b \rightarrow \infty} \left[-\frac{1}{n} \right]_1^b \\
 &= 1 + \lim_{b \rightarrow \infty} \left[-\frac{1}{b} + \frac{1}{1} \right] = 1 + (0 + 1) \\
 \sum_{n=1}^{\infty} \frac{1}{n^2} &< 2, \text{ since the sum is less than 2, the series converges.}
 \end{aligned}$$

Q.N.4) Find the polar equation of the circle $(x+2)^2 + y^2 = 4$

Solution:

$$= (x+2)^2 + y^2 = 4$$

$$= x^2 + 4x + 4 + y^2$$

$$r^2 + 4r \cos \theta = 0$$

$$r^2 = -4r \cos \theta$$

$$r = -4\cos\theta \text{ (Ans)}$$

Q.N.5) Find the area of the parallelogram where vertices are A(0,0), B(7,3), C(9,8) and D(2,5).

Solution:

$$|\vec{AB}| = -\vec{i} + \vec{j}$$

$$|\vec{AD}| = \vec{i} + \vec{j}$$

$$\Rightarrow |\vec{AB}| \times |\vec{AD}| = \begin{vmatrix} i & j \\ -1 & 1 \\ 1 & 1 \end{vmatrix}$$

$$= (-i+j) \times (-i+j)$$

$$= |1 + 1| = |2| = 2$$

Q.N.6) Evaluate the integral

$$\begin{aligned} &= \int_t^{2t} \int_0^t (\sin x + \cos x) dx dy \\ &= \int_t^{2t} [-\cos x + \sin x]_0^t dy \\ &= \int_t^{2t} [-\cos t + \cos 0 + \sin t - \sin 0] dy \\ &= \int_t^{2t} [-\cos t + \sin t + 1] dy \\ &= [-y \cos t + y \sin t + y]_t^{2t} \\ &= -t \cos t + t \sin t + t \end{aligned}$$

Which is the required equation

Q.N.7) Evaluate the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - xy}{\sqrt{x} - \sqrt{y}}$$

Solution:

$$= \lim_{(x,y) \rightarrow (0,0)} \frac{x(x-y)}{\sqrt{x} - \sqrt{y}}$$

$$= \lim_{(x,y) \rightarrow (0,0)} \frac{x(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})}{(\sqrt{x} - \sqrt{y})}$$

$$= \lim_{(x,y) \rightarrow (0,0)} x(\sqrt{x} + \sqrt{y})$$

$$= 0$$

Q.N.8) Find $\left(\frac{\partial \omega}{\partial x}\right) y, z$ if $\omega = x^2 + y - z + \sin t$ and $x + y = t$.

Solution:

$$\frac{\partial \omega}{\partial x} = 2x + \frac{\partial \sin t}{\partial t} X \frac{\partial t}{\partial x}$$

$$2x + \cos t X 1 = 2x + \cos t (x + y)$$

Q.N.9) Solve the partial differential equation $p + q = x$

Solution:

$$p + q = x$$

$$\frac{dx}{1} = \frac{dy}{1} = \frac{dz}{1}$$

$$\Rightarrow \frac{dx}{1} = \frac{dy}{1}$$

$$\Rightarrow x = y - C_1$$

$$\Rightarrow C_1 = x - y \dots \dots \dots (i)$$

$$\Rightarrow \frac{dx}{1} = \frac{dz}{x}$$

$$\Rightarrow x \cdot dx = dz$$

$$\Rightarrow \frac{x^2}{2} = z + C_2$$

$$C_2 = \frac{x^2}{2} - z \dots\dots\dots (ii)$$

$$(C_1, C_2) = (x - y, \frac{x^2}{2} - z)$$

Q.N.10) Find the general integral of the linear partial differential equation

$$z(xp - yq) = y^2 - x^2$$

Solution:

$$z(xp - yq) = y^2 - x^2$$

comparing to the $Pp + Qq = R$

$$P = zx, Q = -y, R = y^2 - x^2$$

$$\frac{dx}{zx} = \frac{dy}{-zy} = \frac{dz}{y^2 - x^2}$$

$$\frac{dx}{x} = \frac{dy}{-xy}$$

$$\log x = -\log y + C_1$$

$$C_1 = \log x + \log y$$

$$\log C_1 = \log xy$$

$$C_1 = xy \dots\dots\dots (i)$$

$$\text{Now, } \frac{dx + dy}{zx - zy} = \frac{dz}{y^2 - x^2}$$

$$\frac{dx + dy}{z(x - y)} = \frac{dz}{(x - y)(x + y)}$$

$$\frac{d(x+y)}{z} = -\frac{dz}{x+y}$$

$$(x+y)d(x+y) = -zdz$$

$$\frac{(x+y)^2}{2} = -\frac{z^2}{2} + C_2$$

$$(C_1, C_2) = 0$$

$$\left(xy, \frac{(x+y)^2}{2} + \frac{z^2}{2} \right) = 0$$

Group B(5X4=20)

Q.N 11) State and prove Rolle's Theorem.

(Rolle's Theorem) Suppose $f(x)$ is continuous on $[a, b]$ and differentiable on (a, b) . If $f(a) = 0 = f(b)$, then there exists a point $c \in (a, b)$ such that $f'(c) = 0$.

Proof. Suppose $f(x)$ is continuous on $[a, b]$, differentiable on (a, b) and $f(a) = 0 = f(b)$. We will prove the theorem using two cases. First, suppose that $f(x) > 0$ for some $x \in (a, b)$. Since $f(x)$ is continuous on $[a, b]$, there exists a point $c \in [a, b]$ for which $f(c)$ is the maximum value of f on $[a, b]$. Furthermore, $f(c) > 0$ implies $c = a$ and $c = b$, so $c \in (a, b)$ and so $f'(c) = 0$ because $f(x)$ is differentiable on (a, b) .

Now suppose $f(x) \leq 0$ for all $x \in (a, b)$. Then either $f(x) = 0$ for all $x \in (a, b)$ in which case $f'(x) = 0$ for all $x \in (a, b)$, or else $f(x) < 0$ for some $x \in (a, b)$. Since $f(x)$

is continuous on $[a, b]$, we know that there is a point $c \in [a, b]$ for which $f(c)$ is the minimum value of $f(x)$ on $[a, b]$. Since $f(c)$ is the minimum on $[a, b]$ and $f(x) < 0$ for some $x \in (a, b)$, $f(c) < 0$. Consequently, $c = a$ and $c = b$, so $c \in (a, b)$ and therefore $f'(c) = 0$ because $f(x)$ is differentiable on (a, b) . This proves the theorem.

Q.N.12) Find the length of the cardioid $r=1+\cos\theta$.

Q.N.13) Define unit tangent vector of a differentiable curve. Find the unit tangent vector of the curve

$$\mathbf{r}(t) = (\cos t + t \sin t) \mathbf{i} + (\sin t - t \cos t) \mathbf{j} \quad t > 0$$

Unit tangent vector of a differentiable curve $\mathbf{r}(t)$ is

$$\mathbf{T} = \frac{d\mathbf{r}}{ds} = \frac{\frac{d\mathbf{r}}{dt}}{\frac{ds}{dt}} = \frac{\mathbf{v}}{|\mathbf{v}|}$$

Solution:

here,

$$(\cos t + t \sin t) \mathbf{i} + (\sin t - t \cos t) \mathbf{j}, t > 0$$

we have,

$$\mathbf{V} = \frac{d\mathbf{r}}{dt} = (-\sin t + \sin t + t \cos t) \mathbf{i} + (\cos t + t \sin t - \cos t) \mathbf{j}$$

$$|\mathbf{V}| = \sqrt{(t \cos t)^2 + (t \sin t)^2} = t$$

$$\mathbf{T} = \frac{\mathbf{v}}{|\mathbf{v}|}$$

$$= (\cos t) \mathbf{i} + (\sin t) \mathbf{j}$$

Q.N.14) what do you mean by critical point of a function $f(x,y)$ in a region? Find local extreme values of the function $f(x,y)=xy-x^2-y^2-2x-2y+4$.

Solution:

An interior point of the domain of a function f where f' is 0 or undefined is called point of function f .

$$f(x, y) = xy - x^2 - y^2 - 2x - 2y + 4$$

$$\text{Here, } f_x = 4 - 2x - 2$$

$$f_y = x - 2y - 2$$

For critical points

$$f_x = 0$$

$$f_y = 0$$

$$\text{ie; } -2x + y = 0$$

$$\text{or; } x - 2y - 2 = 0$$

$$\text{or; } x = -2, y = -2$$

Again,

$$f_{xx} = -2 < 0$$

$$f_{yy} = -2 < 0$$

$$f_{xy} = 1$$

$$\therefore f_{xx} \cdot f_{yy} - f_{xy}^2$$

$$= -2 \cdot -2 - (1)^2$$

$$= 3, \text{ which is } > 0$$

\therefore the function $f(x, y)$ having local maximum is point $(-2, -2)$

$$F_{\max} = -2X - 2 - (-2)^2 - (-2)^2 - 2X - 2 - 2X - 2$$

$$= 8$$

$$\text{critical point } f(x) = 0$$

$$f(y) = 0$$

$$f(x, y) = xy$$

$$f(x) = y$$

$$f(y) = x$$

$$\text{For critical points } f(x) = 0, f(y) = 0$$

$$y = 0$$

$$x = 0$$

$$\therefore \text{critical point is } (0, 0)$$

Again

$$f_{xx} = 0$$

$$f_{yy} = 0$$

$$\therefore f_{xx} \cdot f_{yy} - (f_{xy})^2 = 0 \cdot 0 - (1)^2 = -1$$

$$\therefore f \text{ has local minimum at points } (0, 0)$$

Q.N.15) Find a particular integral of the equation:

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial y^2} = 2y - x^2$$

Solution:

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial y^2} = 2y - x^2$$

$$\text{ie; } (D^2 - D'^2)z = 2y - x^2$$

$$PI = \frac{1}{D^2 - D'^2} (2y - x^2)$$

$$= \frac{\left[1 - \left(\frac{D'}{D}\right)^2\right]^{-1}}{D^2} (2y - x^2)$$

$$= \frac{1}{D^2} \left[2y - x^2 + \frac{D'^2}{D^2} (2y - x^2)\right]$$

$$= \frac{1}{D^2} \left[2y - x^2 + \frac{2yD'}{D^2} - \frac{D'}{D^2} x^2\right]$$

$$= \frac{2y}{D^2} - \frac{x^2}{D^2} + \frac{2yD'}{D^4} - \frac{D'}{D^4} x^2$$

$$= \frac{2x^2y}{2} - \frac{x^4}{12} + \frac{2x^4}{24} - 0$$

$$= x^2y - \frac{x^4}{12} + \frac{x^4}{12}$$

$$= x^2y$$

Group C

Q.N.16) Graph the function $y = x^{4/3} - 4x^{1/3}$

Solution

$$\text{Given equation is } y = x^{4/3} - 4x^{1/3}$$

The first derivative

$$f'(x) = \frac{d}{dx} \left(x^{4/3} - 4x^{1/3} \right) = 4x^{1/3}$$

$$= \frac{4}{3}x^{\frac{1}{3}} - \frac{4}{3}x^{-\frac{2}{3}}$$

$$= \frac{4}{3}x^{-\frac{2}{3}}(x-1)$$

Solving $f'(x) = 0$ we get

$x = 0$ and $x = 1$ which is critical point

Now

$$f''(x) = \frac{4}{9}x^{-\frac{2}{3}} + \frac{8}{9}x^{-\frac{5}{2}}$$

$$= \frac{4}{9}x^{-\frac{5}{2}}(x+2)$$

Solving $f''(x) = 0$ we get

$x=0$ and $x=-2$ which is the required point of inflection





The sign pattern of f' is:

Sign of $\frac{4}{3}x^{\frac{2}{3}}$	+	+	+
Sign of $(x-1)$	-	-	-
Sign of $f'(x) = \frac{4}{3}x^{\frac{2}{3}}$	-	-	+
Change in f	↘	↘	↗

For concavity

Sign of $\frac{4}{9}x^{\frac{5}{3}}$:	+	+	+
Sign of $(x+2)$:	-	+	+
Sign of $\frac{4}{9}x^{\frac{5}{3}}$	-	+	+
	Concave down	Concave up	Concave up

Summary

concave down	Concave up	Concave up	concave up
			




Fig:- Graph (Q.N.16)

Q.N.17) What do you mean by Taylor's Polynomial of order n? Obtain Taylor's polynomial and Taylor's series generated by the function $f(x) = \cos x$ at $x=0$.

Solution:

The functions & its derivatives are

$$f(x) = \cos x \quad f(0) = 1$$

$$f'(x) = -\sin x \quad f'(0) = 0$$

$$f''(x) = -\cos x \quad f''(0) = -1$$

$$f'''(x) = \sin x \quad f'''(0) = 0$$

$$f^{2k}(0) = (-1)^k$$

The series has only even power term for $n=2k$, Taylor's representation is given as:

$$\begin{aligned} \cos x &= f(0) + f'(0)(x-0) + \frac{f''(0)}{2!}(x-0)^2 + \frac{f'''(0)}{3!}(x-0)^3 \\ &\quad + \frac{f^{(4)}(0)}{4!}(x-0)^4 + \dots + \frac{f^{2k}(0)(x-0)^{2k}}{(2k)!} + R_{2k}(x) \end{aligned}$$

$$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + \frac{(-1)^k (x)^{2k}}{(2k)!} + R_{2k}(x)$$

[Note: even term are 0 since every $\sin x = \sin(0) = 0$]

Because the derivative of the cosine have absolute values less than or equal to 1, the remainder estimation theorem with $M=1$ gives

$$|R_{2k}(x)| \leq 1 \cdot \frac{|x|^{2k+1}}{(2k+1)!}$$

For every value of x , $R_{2k} \rightarrow 0$ as $k \rightarrow \infty$. Therefore, the series converges to $\cos x$ for every value of x . Thus,

$$\cos x = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

Q.N.18) Find the volume of the region D enclosed by the surfaces $z = x^2 + 3y^2$ and $z = 8 - x^2 - y^2$.

Solution:

$$z = x^2 + 3y^2 \dots \dots \dots (i)$$

$$z = 8 - x^2 - y^2 \dots \dots \dots (ii)$$

From (i) & (ii)

$$x^2 + 3y^2 = 8 - x^2 - y^2$$

$$\text{or; } 2x^2 + 4y^2 = 8$$

$$\text{or; } 2y^2 = 4 - x^2$$

$$\text{or; } x^2 + 2y^2 - 4 \dots \dots \dots (iii)$$

$$\text{or; } y = \pm \sqrt{\frac{(4 - x^2)}{2}}$$

$$\text{i.e; } -\sqrt{\frac{4-x^2}{2}} \leq \sqrt{\frac{4-x^2}{2}}$$

Now , to find the points for x-axis.

$$y = 0$$

$$\sqrt{\frac{4 - x^2}{2}} = 0$$

$$\therefore x = \pm 2. \quad \text{i.e; } -2 \leq x \leq 2.$$

The points of x-axis are (2,0,0) & (-2,0,0)

Then x² plane will be (2,0,4) & (-2,0,4)

$$\text{volum}(v) = \int_{-2}^2 \int_{-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} \int_{x^2+3y^2}^{8-x^2-y^2} dz dy dx$$

$$\int_{-2}^2 \int_{-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} (8 - 2x^2 - 4y^2) dy dx$$

$$\int_{-2}^2 \left[8y - 2x^2y - \frac{4y^3}{3} \right]_{-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} dx$$

$$\left[\int_{-2}^2 8 \sqrt{\frac{(4-x^2)}{2}} - 2x^2 \sqrt{\frac{(4-x^2)}{2}} - 4 \left(\sqrt{\frac{(4-x^2)}{2}} \right)^3 + 8 \sqrt{\frac{(4-x^2)}{2}} \right. \\ \left. - 2x^2 \sqrt{\frac{(4-x^2)}{2}} - \frac{4}{3} \left(\sqrt{\frac{(4-x^2)}{2}} \right)^3 \right] dx$$

$$= \left[\int_{-2}^2 \left[16 \sqrt{\frac{(4-x^2)}{2}} - 4x^2 \sqrt{\frac{(4-x^2)}{2}} - \frac{8}{3} \left(\sqrt{\frac{(4-x^2)}{2}} \right)^3 \right] dx \right]$$

$$= \int_0^2 \frac{8}{3\sqrt{2}} (4-x^2)^{\frac{3}{2}} dx$$

Put $x = 2\sin\theta$

$$dx = 2\cos\theta d\theta$$

$$= \frac{16}{3\sqrt{2}} \int_0^{\frac{\pi}{2}} (4 - 4\sin^2 \theta)^{\frac{3}{2}} 2\cos\theta d\theta$$

$$\frac{32}{3\sqrt{2}} \int_0^{\frac{\pi}{2}} (4\cos^2 \theta)^{\frac{3}{2}} \cos\theta d\theta$$

$$= \frac{32}{3\sqrt{2}} \int_0^{\frac{\pi}{2}} 8\cos^4 \theta d\theta$$

$$= \frac{32}{3\sqrt{2}} \int_0^{\frac{\pi}{2}} 2(2\cos^2 \theta)^2 d\theta$$

$$\frac{64}{3\sqrt{2}} \int_0^{\frac{\pi}{2}} (1 + \cos \theta)^2 d\theta$$

Therefore volume = $8\sqrt{2}\pi$

$$\therefore \text{volume}(v) = 8\sqrt{2}$$

Q.N.19) Obtain the absolute maximum and minimum values of the function $f(x,y)=2+2x+2y-x^2-y^2$ on the triangular plate in the first quadrant bounded by lines $x=0, y=0, y=9-x$.

Solution: Since f is differentiable, the only places where can assume these values are points inside the triangle where $f_x=f_y=0$ and points on the boundary .

a) **Interior points** . For these we have

$$f_x=2-2x=0, f_y=2-2y=0,$$

yielding the single point $(x,y)=(1,1)$. The value of f there is $f(1,1)=4$

b) **Boundary points**. We take the triangle one side at a time.

i) **On the segment OA, $y=0$** . The function

$$f(x,y)=f(x,0)=2+2x-x^2$$

may now be regarded as function of x defined on the closed interval

$0 \leq x \leq 9$. Its extreme values may occur at the endpoints

$x=0$ where $f(0,0)=2$

$x=9$ where $f(9,0)=2+18-81= -61$

and at the interior points where $f'(x,0)=2-2x=0$. The only interior point

where $f'(x,0)=0$ is $x=1$, where

$$f(x,0)=f(1,0)=3$$

ii) **On the segment OB, $x=0$** and

$$f(x,y)=f(0,y)=2+2y-y^2$$

we know from the symmetry of f in x and y and from the analysis we just carried out that the candidates on this segment are

$$f(0,0)=2, \quad f(0,9)= -61, \quad f(0,1)=3.$$

iii) we have already accounted for the values of f at the endpoints of AB, so we need only

$$f(x,y)=2+2x+2(9-x)-x^2-(9-x)^2= -61 + 18x - 2x^2$$

setting $f'(x, 9-x) = 18 - 4x = 0$ gives

$$x = \frac{18}{4} = \frac{9}{2}.$$

At this value of x ,

$$y = 9 - \frac{9}{2} = \frac{9}{2} \text{ and } f(x, y) = f\left(\frac{9}{2}, \frac{9}{2}\right) = -\frac{41}{2}.$$

Summary, We list all the candidates: 4, 2, -61, 3, -(41/2). The maximum is 4, which f assumes at (1, 1). The minimum is -61, which f assumes at (0, 9) and (9, 0).

OR

Q.N.19) Evaluate the integral

$$\int_0^1 \int_0^{3-3x} \int_0^{3-3x-y} dz dy dx$$

Solution:

$$\begin{aligned} &= \int_0^1 \int_0^{3-3x} (3-3x-y) dy dx \\ &= \int_0^1 \left[3y - 3xy - \frac{y^2}{2} \right]_0^{3-3x} dx \\ &= \int_0^1 3(3-3x) - 3x(3-3x) - \frac{(3-3x)^2}{2} dx \\ &= \int_0^1 9 - 9x - 9x + 9x^2 - \frac{9 - 18x + 9x^2}{2} dx \\ &= \frac{1}{2} \int_0^1 18 - 18x - 18x + 18x^2 - 9 + 18x - 9x^2 dx \\ &= \frac{1}{2} \int_0^1 9x^2 - 18x + 9 dx \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} [3x^3 - 9x^2 + 9x]_0^1 \\
 &= \frac{1}{2} [3 \times 1 - 9 \times 1^2 + 9 \times 1] \\
 &= \frac{3}{2}
 \end{aligned}$$

Q.N.20) Show the solution of the wave equation.

Solution:

The wave equation $\frac{d^2u}{dt^2} = \frac{C^2 d^2u}{dx^2}$

where $C^2 = \frac{T}{\rho}$

The solution of equation(i) can be obtained by introducing two independent variables v and z defined by $v = x + ct$ and $z = x - ct$

Differentiating v w.r.t x

$$\begin{aligned}
 \frac{dv}{dx} &= \frac{du}{dv} \cdot \frac{dv}{dx} + \frac{du}{dz} \cdot \frac{dz}{dx} \\
 &= \frac{du}{dv} + \frac{du}{dz}
 \end{aligned}$$

Again differentiate w.r. to x

$$\begin{aligned}
 \frac{d^2}{dv} \left(\frac{du}{dv} \cdot \frac{du}{dz} \right) \frac{dv}{dx} + \frac{d}{dz} \left(\frac{du}{dv} + \frac{du}{dz} \right) \frac{dz}{dx} \\
 = \frac{d^2u}{dv^2} + \frac{d^2u}{dv^2 z} + \frac{d^2u}{dz dv} + \frac{d^2u}{dz^2} \\
 \frac{d^2u}{dv^2} + \frac{z d^2u}{dv dz} + \frac{d^2u}{dz^2}
 \end{aligned}$$

Again diff u w.r.to t .

$$\frac{du}{dt} = \frac{du}{dv} \cdot \frac{dv}{dt} + \frac{du}{dz} \cdot \frac{dz}{dt}$$

$$= \frac{du}{dv} \cdot c + \frac{du}{dz} (-c)$$

$$= \frac{cdv}{dv} - \frac{cdv}{dz}$$

$$\frac{d^2u}{dt^2} = c \left[\frac{du}{dv} \left(\frac{du}{dv} - \frac{du}{dz} \right) \frac{dv}{dt} + \frac{du}{dz} \left(\frac{du}{dv} - \frac{du}{dz} \right) \frac{dz}{dt} \right]$$

$$c^2 \left[\frac{d^2u}{dv^2} - \frac{d^2u}{dv^2 dz} - \frac{d^2u}{dz dv} + \frac{d^2u}{dz^2} \right]$$

$$= C^2 \left[\frac{d^2u}{dv^2} - \frac{2d^2u}{dv dz} + \frac{d^2u}{dz^2} \right]$$

Inserting these value in eqn (i)

We get,

$$C^2 \left[\frac{d^2u}{dv^2} - \frac{2d^2u}{dv dz} + \frac{d^2u}{dz^2} \right] = C^2 \left(\frac{d^2u}{dv^2} + \frac{2d^2u}{dv dz} + \frac{d^2u}{dz^2} \right)$$

$$\frac{4d^2u}{dv dz} = 0$$

$$\text{or; } \frac{d^2u}{dv dz} = 0 \dots \dots (ii)$$

To get solution integrating partially w.r. to z .

$$\frac{du}{dv} = \gamma(v)$$

Integrating w.r to v

$$u = \int \gamma(v) dv + \varphi(z)$$

$$u(x, t) = \phi(v) + \varphi(z)$$

$$= \phi(x + ct)$$

$$+ \phi(x$$

$- ct)$ is known as *D'Alembert's solution of wave equation*.

OR

Find a particular integral of the equation $(D^2 - d')z = A \cos(lx + my)$ where A, l, m are constants.

solution;

given equation is $(D^2 - D')z = A \cos(lx + my)$

Given equation is $(D^2 - D')z = A \cos(lx + my) \dots \dots (i)$

Let

$z = C_1 \cos(lx + my) + C_2 \sin(lx + my) \dots \dots (ii)$ be the solution of (i) where C_1 and C_2 are constant to be determined. then from equation (i) and (ii)

$$(D^2 - D') [C_1 \cos(lx + my) + C_2 \sin(lx + my)] = A \cos(lx + my)$$

$$\text{or; } D^2 [C_1 \cos(lx + my) + C_2 \sin(lx + my)] - D' (C_1 \cos(lx + my) + C_2 \sin(lx + my)) = A \cos(lx + my)$$

$$A_1 = D^2 [C_1 \cos(lx + my) + C_2 \sin(lx + my)]$$

$$= D [-C_1 \sin(lx + my) \cdot l + C_2 \cos(lx + my) \cdot l]$$

$$- C_1 l^2 \cos(lx + my) - C_2 l^2 \sin(lx + my)$$

$$B_1 = D' [C_1 \cos(lx + my) + C_2 \sin(lx + my)]$$

$$= -C_1 m \sin(lx + my) + C_2 m \cos(lx + my)$$

$$= C_2 m \cos(lx + my) - C_1 m \sin(lx + my)$$

solution equation (iii) becomes

$$A_1 - B_1 = A \cos(lx + my)$$

$$\Rightarrow -C_1 l^2 \cos(lx + my) - C_2 l^2 \sin(lx + my) - C_2 m \cos(lx + my) + C_1 m \sin(lx + my) = A \cos(lx + my)$$

$$\Rightarrow (-C_1 l^2 - C_2 m) \cos(lx + my) + (C_1 m - C_2 l^2) \sin(lx + my) = A \cos(lx + my)$$

Comparing the coefficient of like terms on the both side

$$-C_1 l^2 - C_2 m = A \dots \dots \dots (iv)$$

$$C_1 m - C_2 l^2 = 0 \dots \dots \dots (v)$$

From equation(iv)

$$C_2 = -\frac{C_1 l^2 - A}{m}$$

putting C_2 on equatin (v)

$$C_1 m + C_1 l^4 + A l^2 = 0$$

$$\Rightarrow C_1 = -\frac{A l^2}{m^2 + l^4}$$

$$\text{Now, } C_2 \text{ becomes } C_2 = -\frac{A m}{m^2 + l^4}$$

Now, the solution on equation (iii) becomes

$$z = -\frac{A l^2}{m^2 + l^4} \cos(lx + my) - \frac{A m}{l^4 + m^2} \sin(lx + my)$$